

Carboplatin and paclitaxel are to be given only after veliparib/placebo dosing on Day -2 and Day -1 are confirmed.

Table 1. Treatment Schema for Each Cycle

Week 1							
Days	-2	-1	1	2	3	4	5
Veliparib/Placebo	Twice a day	Twice a day	Twice a day	Twice a day	Twice a day	Twice a day	Twice a day
Paclitaxel			Once				
Carboplatin			Once				
Week 2							
	6	7	8	9	10	11	12
Veliparib/Placebo							
Paclitaxel			Once				
Carboplatin							
Week 3							
	13	14	15	16	17	18	19
Veliparib/Placebo							
Paclitaxel			Once				
Carboplatin							

Subjects will continue to receive veliparib/placebo in combination with carboplatin/paclitaxel until unacceptable toxicity occurs or radiographic progression occurs. Subjects who experience toxicities due to carboplatin/paclitaxel or veliparib may require a delay in dosing or dose modification as described in Section 5.7.

Subjects who discontinue carboplatin and paclitaxel due to toxicity and who have not progressed are eligible to receive single-agent, blinded veliparib/placebo starting at 300 mg BID. If the subject tolerates 300 mg BID for 2 weeks, veliparib/placebo may be increased to 400 mg BID at the investigator's discretion. Dosing with veliparib/placebo will begin on Day -2 and continue through Day 19 of a 21-day cycle. Subjects will continue to follow the schedule of assessments as outlined in Table 2.